

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 235/52, 237/22, C07D 207/16, C07C 323/02, 231/02, A61K 31/22, 31/165	A1	(11) International Publication Number: WO 98/56756 (43) International Publication Date: 17 December 1998 (17.12.98)
(21) International Application Number: PCT/KR97/00156 (22) International Filing Date: 20 August 1997 (20.08.97) (30) Priority Data: 1997/24307 12 June 1997 (12.06.97) KR (71) Applicant: CHEIL JEDANG CORPORATION [KR/KR]; Cheil Building, 500, 5-ga, namdaemun-ro, Chung-ku, Seoul 100-095 (KR). (72) Inventors: RHEE, Chung, Keun; Woosunggreen Apt. 106-1303, San2-106, Dapshippli 4-dong, Dongdaemun-ku, Seoul 130-034 (KR). XIANG, Myung, Xik; Injunprince Apt. 1004-1002, Youbangdong, 1007-1, Yongin-shi, Kyungkee-do 449-800 (KR). SUH, Byoung, Chool; Munjungshiyong Apt. 7-802, Munjungdong, Songpa-ku, Seoul 138-200 (KR). LEE, Kwang, Hyuk; 185, Yatopdong, Boondang-ku, Sungnamshi, Kyungkeedo 463-070 (KR). LEE, Youn, Ha; Sambolife Apt. 702, 301, Kimrang- jang-dong, Yongin-shi, Kyungkee-do 449-800 (KR). BHANG, Won, Young; 36-6, Jungja-dong, Bundang-ku, Sungnamshi, Kyungkee-do 463-010 (KR). KIM, Young, Gi; Dunchundong 52-7, 202, Kangdong-ku, Seoul 134-062 (KR).		(74) Agents: CHOI, Hak, Hyun et al.; 6th floor, Peeres Building, 222, 3-ka, Chungjung-ro, Seodaemun-ku, Seoul 120-013 (KR). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: CATECHOL AMINO ACID DERIVATIVES AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM (57) Abstract <p>A phosphodiesterase IV inhibiting catechol derivative of general formula (I) or a pharmaceutically acceptable salt thereof in which R1, R2, R3, R4, R5 and W are as defined herein. In addition, a process for producing the compound of general formula (I) and a pharmaceutical composition containing pharmaceutically effective amount of the compound of general formula (I) are disclosed.</p> <div data-bbox="795 1155 1218 1333"><p>(I)</p></div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

CATECHOL AMINO ACID DERIVATIVES AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM

5 **FIELD OF THE INVENTION**

 The present invention relates to novel catechol amino acid derivatives which inhibit the enzymatic activity of phosphodiesterase IV or tumor necrosis factor. These compounds may be useful in prevention or treatment of bronchial
10 asthma, arthritis, bronchitis, chronic atretic airway, psoriasis, allergic rhinitis, dermatitis, AIDS, Crohn's disease, septicemia, septic shock, other inflammatory diseases such as cachexia, TNF related diseases, etc. Also, the present invention relates to a method for producing the said compounds and a
15 pharmaceutical composition containing the said compounds.

15 **BACKGROUND OF THE INVENTION**

 Phosphodiesterase IV is an enzyme that specifically hydrolyzes cAMP (adenosine 3',5'-cyclic monophosphate) into inactive adenosine 3',5'-mono-
20 phosphate. The cAMP has been shown to be a second messenger mediating the cellular responses to external stimuli and to act as relaxing or contradicting bronchial muscles.

 The inhibition of phosphodiesterase IV leads to the prevention of
25 broncospasm by maintaining the concentration of cAMP and also induces an anti-inflammation. Therefore, compounds that inhibit phosphodiesterase IV should be effective in treating asthma and the like diseases.


 It is known that tumor necrosis factor (TNF) is implicated in
30 infectious disease such as cachexia and autoimmune disease. Also TNF appears

to act as a primary mediator for inflammatory reaction such as septicemia and septic shock.

Therefore, it is expected that compounds with the inhibitory activity against phosphodiesterase IV or TNF will be pharmaceutically valuable and there is always a need to develop new compounds which inhibit phosphodiesterase IV and TNF.

SUMMARY OF THE INVENTION

In one aspect, the present invention provides a compound of the general formula I:



Formula I

or a pharmaceutically acceptable salt or solvate thereof in which

R1 represents methyl, ethyl, difluoromethyl or trifluoromethyl; **R2** represents C₁-C₇ alkyl optionally substituted with nitro or halogen, C₃-C₇ cycloalkane optionally substituted with nitro or halogen, phenyl optionally substituted with nitro or halogen, C₂-C₅ heteroaryl including N, O or S optionally substituted with nitro or halogen; **R3** represents hydrogen, hydroxy, C₁-C₆ alkyl or phenyl substituted with C₁-C₄ alkyl, C₁-C₄ alkoxy, nitro or halogen; **R4** represents hydrogen, trifluoromethyl, C₁-C₆ alkyl, -(CH₂)_nCO₂H, -(CH₂)_nCONH₂, -(CH)_n phenyl, -(CH)_n phenylalcohol, -(CH₂)_n indole, -(CH₂)_nimidazole, -(CH₂)_xOH,

5 $-(CH_2)_xSH$, $-(CH)_xS$ C_1-C_6 alkyl, $-(CH_2)_xNHC(NH)NH_2$ or $-(CH_2)_xNH$ in which x is 1 to 5 and n is 0 to 5; or R_3 and R_4 are together taken with C_1-C_4 to form a ring; R_5 represents NR_6R_7 , $-OR_8$ or $-OCH(R_8)OCOR_9$ in which R_6 and R_7 are each independently hydrogen, C_1-C_6 alkyl or phenyl substituted with C_1-C_4 alkoxy, nitro or halogen and R_8 and R_9 are each independently C_1-C_8 alkyl, C_3-C_{11} cycloalkyl substituted with halogen, nitro, hydroxy, C_1-C_4 alkyl or C_1-C_4 alkoxy or phenyl substituted with halogen, nitro, hydroxy, C_1-C_4 alkyl or C_1-C_4 alkoxy; and W represents oxygen or sulfur.

10 In another aspect, the present invention provides a process for producing the above compound of the general formula I.

15 In still another aspect, the present invention provides a pharmaceutical composition comprising pharmaceutically effective amount of the present compound of the general formula I or pharmaceutically acceptable salt or solvate thereof and a pharmaceutically acceptable carrier.

DETAILED DESCRIPTION OF THE INVENTION


20 The preferred compounds of the present invention are those wherein R_1 is methyl; R_2 is cyclopentyl or norbornyl; R_3 is hydrogen; R_4 is hydrogen, C_1-C_4 alkyl or CH_3SH ; or R_3 and R_4 are together taken with C_4 to form a ring; R_5 is NR_6R_7 , $-OR_8$ or $-OCH(R_8)OCO_2R_9$ in which R_6 and R_7 are each independently hydrogen or C_1-C_4 alkyl and R_8 and R_9 are each
25 independently C_1-C_4 alkyl; and W is oxygen.

The more preferred compounds of the present invention are as follows:

30 ethyl-2-(3-cyclopentyloxy-4-methoxyphenylcarboxylamido)-3-sulfamyl-(2S)-propanoate;

1N-carbamoylmethyl-3-cyclopentyloxy-4-methoxybenzylamide; and
ethyl-2-(3-cyclopentyloxy-4-methoxyphenylcarboxylamido)acetate.


5 The process for producing the compound of the general formula I according to the present invention comprises reacting a compound of the general formula II:



15

Formula II

in which R1, R2 and W represent the same as defined above, with a compound of the general formula III:




25

Formula III

in which R3, R4 and R5 are the same as defined above, to produce the
30 compound of the general formula I.

The above compound II was obtained by known method (J. Med. Chem. 1994, 37, 1696). This method is represented by the following reaction scheme:

5



The above compound III is commercially available.

For use in medicine, the compounds of the present invention are usually administered as a standard pharmaceutical composition. Therefore, the present invention provides in a further aspect pharmaceutical compositions comprising a novel compound of formula I or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

The compounds of formula I may be administered by any convenient method, for example by oral, parenteral, buccal, sublingual, nasal, rectal or transdermal administration and the pharmaceutical compositions adapted accordingly.

The compounds of formula I and their pharmaceutically acceptable salts which are active when given orally can be formulated as liquids, for example syrups, suspensions or emulsions, tablets, capsules and lozenges.

A liquid formulation will generally consist of a suspension or solution of the compound or physiologically acceptable salt in a suitable liquid carrier, for example an aqueous solvent such as water, ethanol or glycerine, or a

non-aqueous solvent, such as polyethylene glycol or an oil. The formulation may also contain a suspending agent, preservative, flavouring or colouring agent.

5 A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations. Examples of such carriers include magnesium stearate, starch, lactose, sucrose and cellulose.

10 A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatin capsule; alternatively, a dispersion or suspension can be prepared using any suitable pharmaceutical carrier(s), for example aqueous gums, celluloses,
15 silicates or oils and the dispersion or suspension then filled into a soft gelatin capsule.

 Typical parenteral compositions consist of a solution or suspension of the compound or physiologically acceptable salt in a sterile aqueous carrier or
20 parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilized and then reconstituted with a suitable solvent prior to administration.

25 Compositions for nasal administration may conveniently be formulated as aerosols, drops, gels and powders. Aerosol formulations typically comprise a solution or fine suspension of the active substance in a physiologically acceptable aqueous or non-aqueous solvent and are usually presented in single or multidose quantities in sterile form in a sealed container, which can take
30 the form of a cartridge or refill for use with an atomising device.

Alternatively the sealed container may be a unitary dispensing device such as a single dose nasal inhaler or an aerosol dispenser fitted with a metering valve which is intended for disposal once the contents of the container have been exhausted. Where the dosage form comprises an aerosol dispenser, it will
5 contain a propellant which can be a compressed gas such as compressed air or an organic propellant such as a fluorochlorohydrocarbon. The aerosol dosage forms can also take the form of a pump-atomiser.

Compositions suitable for buccal or sublingual administration include
10 tablets, lozenges and pastilles, wherein the active ingredient is formulated with a carrier such as sugar and acacia, tragacanth, or gelatin and glycerin.

Compositions for rectal administration are conveniently in the form of suppositories containing a conventional suppository base such as cocoa butter.
15

Compositions suitable for transdermal administration include ointments, gels and patches.

Preferably the composition is in unit dose form such as a tablet,
20 capsule or ampoule.

The invention will now be described with reference to the following illustrative Examples.

25 **EXAMPLES**

REFERENCE EXAMPLE 1

3-Cyclopentyloxy-4-methoxybenzylaldehyde

30 50 g of isovanillin was added dropwise to 300 ml of

dimethylformamide and, then, 70 g of anhydrous potassium carbonate and 1.5 g of potassium iodide were added to form a suspension. The suspension was stirred at 65°C for 30 minutes. 63 g of cyclopentyl bromide was added dropwise to the suspension over 1 hour and was stirred at 65°C for 1 day.

5 After cooling to room temperature, the suspension was diluted by adding 1 L of toluene and, then, was washed with 1 N sodium hydroxide (2x500 ml) and distilled water (2x250 ml) in succession. The organic layer was dried over anhydrous sodium sulfate and was concentrated under reduced pressure to obtain 58 g of the title compound (yield: 80%).

10 ¹H NMR(CDCl₃, δ): 9.84(1H,s), 7.42(2H,m), 6.94(1H,D,J=9Hz), 4.87(1H,m), 3.93(3H,s), 2.1-1.6(8H,m)

REFERENCE EXAMPLE 2

3-Cyclopentyloxy-4-methoxybenzoic acid

15 35 g of sulfamic acid was added dropwise to a solution of 58 g of 3-cyclopentyloxy-4-methoxybenzaldehyde in 450 ml of 80% acetic acid. Then, a solution of 30 g of 80% sodium chloride in 450 ml of distilled water was added dropwise to the suspension over 1 hour while maintaining the reaction temperature to 18°C-20°C. The reaction solution was stirred for 1 hour

20 and was diluted by adding dropwise 450 ml of distilled water over 30 minutes. The solution was filtered with distilled water and was dried to obtain 56 g of white solid title compound (yield: 90%).

¹H NMR(CDCl₃, δ): 7.73(1H,dd,J=9,1Hz), 7.24(2H,d,J=1Hz), 6.92(1H,d,J=9Hz),

25 4.84(1H,m), 3.93(3H,s), 2.1-1.6(8H,m)

REFERENCE EXAMPLE 3

3-Cyclopentyloxy-4-methoxybenzoic acid chloride

30 54 g of 3-cyclopentyloxy-4-methoxybenzoic acid was added to 30 ml

of thiocloride and was stirred for 5 hours. 50 ml of toluene was added to the reaction solution and was concentrated under reduced pressure to obtain 58 g of the dense brown title compound (yield: 98%).

¹H NMR(CDCl₃, δ): 7.82(1H,dd,J=9,1Hz), 7.53(1H,d,J=1Hz), 6.92(1H,d,J=9Hz),
5 4.9-4.8(1H,m), 3.87(3H,s), 2.1-1.9(2H,m), 1.9-1.7(4H,m), 1.7-1.5(2H,m)

EXAMPLE 1

Ethyl-2-(3-cyclopentyloxy-4-methoxyphenylcarboxylamido)-3-phenyl-(2S)-propanoate

10

10 ml of pyridine was added dropwise to 400 mg of L-phenylalanine to form a suspension. 500 mg of 3-cyclopentyloxy-4-methoxybenzoic acid chloride (Reference Example 3) was added dropwise to the suspension and was refluxed for 4 hours. The reaction solution was concentrated under
15 reduced pressure, diluted with dichloromethane and washed with 1N hydrochloric acid, saturated sodium bicarbonate and sodium chloride in succession. The organic layer was dried and concentrated under reduced pressure. The solid compound obtained therefrom was recrystallized with toluene to yield 630 mg of the solid thin yellow title compound. m.p.
20 105-107°C

¹H NMR(DMSO-d₆, δ): 7.23(1H,d), 7.20(4H,m), 7.05(2H,m), 6.72(1H,H,d), 6.40(1H,d), 4.90(1H,ddd), 4.70(1H,m), 4.12(2H,q), 3.77(3H,s), 3.12(2H,ddd), 1.76(6H,m), 1.40(2H,m), 1.12(3H,t)

25 EXAMPLE 2

Ethyl-2-(3-cyclopentyloxy-4-methoxyphenylcarboxylamido)-3-sulfanyl-(2S)-propanoate

321 mg of L-cysteine ethylester was reacted with 400 mg of
30 3-cyclopentyloxy-4-methoxybenzoic acid chloride (Reference Example 3) in the

same manner as described in Example 1 to yield 600 mg of the title compound. m.p. 54-56°C

1H NMR(CDCl₃, δ): 7.60(1H,dd), 7.41(1H,d), 7.41(1H,d), 7.38(1H,d), 6.80(1H,d), 5.00(1H,ddd), 4.85(1H,m), 4.25(2H,q), 3.84(3H,s), 3.65(1H,ddd),
5 3.17(1H,ddd), 1.70(8H,m), 1.28(3H,t)

EXAMPLE 3

1N-[1-carbamoyl-2-phenyl-(1S)-ethyl-3-cyclopentyloxy-4-methoxybenzylamide

10 10 ml of ammonium hydroxide was added to a solution of 780 mg of ethyl-2-(3-cyclopentyloxy-4-methoxyphenylcarboxylamido)-3-phenyl-(2S)-propanoate (Example 1) in 50 ml of methanol and was refluxed for 10 hours. After cooling the reaction solution, the solids formed was filtered and recrystallized with toluene to yield 450 mg of the solid white title compound.
15 m.p. 139-140°C

1H NMR(DMSO-d₆, δ): 8.69(1H,d), 7.43(1H,d), 7.25(5H,m), 7.20(1H,dd), 6.99(1H,d), 4.80(1H,m), 4.62(1H,m), 3.79(3H,s), 3.13(2H,ddd), 1.75(8H,m)

EXAMPLE 4

20 1N-carbamoylmethyl-3-cyclopentyloxy-4-methoxybenzylamide

385 mg of glycineamide hydrochloride was reacted with 1.0 g of 3-cyclopentyloxy-4-methoxybenzoic acid chloride (Reference Example 3) in the same manner as described in Example 1 to yield 600 mg of the white title
25 compound. m.p. 131-132°C

1H NMR(CDCl₃, δ): 7.45(1H,d), 7.38(1H,dd), 7.2-7.0(1H,brs), 6.89(1H,d), 6.5-6.3(1H,brs), 5.7-5.5(1H,brs), 5.0-4.8(1H,m), 4.18(2H,d), 3.91(3H,s), 2.1-1.8(4H,m), 1.8-1.6(2H,m)

30 EXAMPLE 5

Ethyl-2-(3-cyclopentyloxy-4-methoxyphenylcarboxylamido)acetate

490 mg of glycine ethylester hydrochloride was reacted with 1.0 g of 3-cyclopentyloxy-4-methoxybenzoic acid chloride (Reference Example 3) in the same manner as described in Example 1 to yield 950 mg of the light yellow title compound. m.p. 73-74°C

¹H NMR(CDCl₃, δ): 7.50(1H,d), 7.41(1H,dd), 6.95(1H,d), 6.7-6.6(1H,brs), 5.9-5.8(1H,m), 4.34(2H,q), 4.30(2H,d), 3.97(3H,s), 2.0-1.6(8H,m), 1.40(3H,t)

EXAMPLE 6**Methyl-2-(3-cyclopentyloxy-4-methoxyphenylcarboxylamido)-4-methyl-(2S)-pentanoate**

785 mg of L-leucine ethylester hydrochloride was reacted with 1.0 g of 3-cyclopentyloxy-4-methoxybenzoic acid chloride (Reference Example 3) in the same manner as described in Example 1 to yield 900 mg of the white title compound. m.p. 123-124°C

¹H NMR(CDCl₃, δ): 7.43(1H,d), 7.29(1H,dd), 6.87(1H,d), 6.43(1H,d), 4.9-4.7(1H,m), 3.90(3H,s), 3.78(3H,s), 2.0-1.6(11H,m), 1.01(3H,d), 0.99(3H,d)

20

EXAMPLE 7**1-(3-cyclopentyloxy-4-methoxybenzoyl)-(2S)-perhydro-2-pyrrolecarboxylamide**

460 mg of L-prolineamide was reacted with 1.0 g of 3-cyclopentyloxy-4-methoxybenzoic acid chloride (Reference Example 3) in the same manner as described in Example 1 to yield 500 mg of the white title compound. m.p. 170-171°C

¹H NMR(CDCl₃, δ): 7.2-7.0(2H,m), 7.0-6.8(1H,brs), 6.89(1H,d), 5.7-5.5(1H,brs), 4.9-4.8(2H,m), 3.90(3H,s), 3.8-3.6(2H,m), 2.6-2.4(1H,m), 2.2-1.6(11H,m)

30

EXAMPLE 8**Methyl-1-(3-cyclopentyl-4-methoxybenzoyl)-(2S)-perhydro-2-pyrrolecarboxyl-ester**

5 670 mg of L-prolinemethylester was reacted with 1.0 g of 3-cyclopentyl-4-methoxybenzoic acid chloride (Reference Example 3) in the same manner as described in Example 1 to yield 500 mg of the liquid colorless title compound.

10 ¹H NMR(CDCl₃, δ): 7.2-7.0(2H,m), 6.78(1H,d), 4.9-4.8(1H,m), 4.7-4.6(1H,m), 3.90(3H,s), 3.78(3H,s), 3.8-3.6(2H,m), 2.2-1.6(12H,m)

EXPERIMENTAL EXAMPLE**Inhibition of phosphodiesterase IV activity**

15

Compounds prepared by Examples 1 to 7 and Rolipram as control were tested on the inhibition of phosphodiesterase IV.

20 Phosphodiesterase IV partially purified from human U937 cells, test compound and 1.0 μM cAMP including 0.01 μM [³H] cAMP were incubated at 30°C for 20 minutes. The PDE reaction to convert cAMP into AMP was completed by boiling the reaction solution for 2 minutes. AMP was converted into adenosine by adding snake venom nucleotidase and incubating the reaction solution at 30°C for 10 minutes. While unhydrolyzed cAMPs were bonded to
25 AG1-X2 resin, the [³H] adenosine in the aqueous solution was quantified by scintillation counting. The results are shown in Table II below, in which the values indicate inhibition(%) of the PDE IV by each test compound.


30

Table I

Test Compound	Example No.							Rolipram
	1	2	3	4	5	6	7	
Inhibition(%)	74.5	38.2	72.4	36.5	74.7	77.9	65.6	59.7

WHAT IS CLAIMED IS:

1. A compound of the general formula I:



Formula I

or a pharmaceutically acceptable salt or solvate thereof in which

R1 represents methyl, ethyl, difluoromethyl or trifluoromethyl; **R2** represents
 15 C₁-C₇ alkyl optionally substituted with nitro or halogen, C₃-C₇ cycloalkane
 optionally substituted with nitro or halogen, phenyl optionally substituted with
 nitro or halogen, C₂-C₃ heteroaryl including N, O or S optionally substituted
 with nitro or halogen; **R3** represents hydrogen, hydroxy, C₁-C₆ alkyl or phenyl
 substituted with C₁-C₄ alkyl, C₁-C₄ alkoxy, nitro or halogen; **R4** represents
 20 hydrogen, trifluoromethyl, C₁-C₆ alkyl, -(CH₂)_nCO₂H, -(CH₂)_nCONH₂, -(CH)_n
 phenyl, -(CH)_n phenylalcohol, -(CH₂)_n indole, -(CH₂)_nimidazole, -(CH₂)_xOH,
 -(CH₂)_xSH, -(CH)_xS C₁-C₆ alkyl, -(CH₂)_xNHC(NH)NH₂ or -(CH₂)_xNH in which
 x is 1 to 5 and n is 0 to 5; or **R3** and **R4** are together taken with C₁-C₄ to
 form a ring; **R5** represents NR₆R₇, -OR₈ or -OCH(R₈)OCOR₉ in which R₆
 25 and R₇ are each independently hydrogen, C₁-C₆ alkyl or phenyl substituted
 with C₁-C₄ alkoxy, nitro or halogen and R₈ and R₉ are each independently
 C₁-C₈ alkyl, C₃-C₁₁ cycloalkyl substituted with halogen, nitro, hydroxy, C₁-C₄
 alkyl or C₁-C₄ alkoxy or phenyl substituted with halogen, nitro, hydroxy, C₁-C₄
 alkyl or C₁-C₄ alkoxy; and **W** represents oxygen or sulfur.


30

2. The compound of claim 1 wherein R1 is methyl; R2 is cyclopentyl or norbornyl; R3 is hydrogen; R4 is hydrogen, C₁-C₄ alkyl or CH₃SH; or R3 and R4 are together taken with C₄ to form a ring; R5 is NR₆R₇, -OR₈ or -OCH(R₈)OCO₂R₉ in which R₆ and R₇ are each independently hydrogen or C₁-C₄ alkyl and R₈ and R₉ are each independently C₁-C₄ alkyl; and W is oxygen.

3. The compound of claim 1 which is in the form of optical isomer or geometrical isomer.


4. The compound of claim 1 which is ethyl-2-(3-cyclopentyloxy-4-methoxyphenylcarboxylamido)-3-phenyl-(2S)-propanoate, ethyl-2-(3-cyclopentyloxy-4-methoxyphenylcarboxylamido)-3-sulfanyl-(2S)-propanoate, 1N-[1-carbamoyl-2-phenyl-(1S)-ethyl-3-cyclopentyloxy-4-methoxybenzylamide, 1N-carbamoylmethyl-3-cyclopentyloxy-4-methoxybenzylamide, ethyl-2-(3-cyclopentyloxy-4-methoxyphenylcarboxylamido)acetate, methyl-2-(3-cyclopentyloxy-4-methoxyphenylcarboxylamido)-4-methyl-(2S)-pentanoate, 1-(3-cyclopentyloxy-4-methoxybenzoyl)-(2S)-perhydro-2-pyrrolicarboxylamide, or methyl-1-(3-cyclopentyl-4-methoxybenzoyl)-(2S)-perhydro-2-pyrrolicarboxylester.

5. A process for producing a compound of the general formula I:




Formula I

or a pharmaceutically acceptable salt or solvate thereof in which **R1** represents methyl, ethyl, difluoromethyl or trifluoromethyl; **R2** represents C_1 - C_7 alkyl optionally substituted with nitro or halogen, C_3 - C_7 cycloalkane optionally substituted with nitro or halogen, phenyl optionally substituted with nitro or halogen, C_2 - C_5 heteroaryl including N, O or S optionally substituted with nitro or halogen; **R3** represents hydrogen, hydroxy, C_1 - C_6 alkyl or phenyl substituted with C_1 - C_4 alkyl, C_1 - C_4 alkoxy, nitro or halogen; **R4** represents hydrogen, trifluoromethyl, C_1 - C_6 alkyl, $-(CH_2)_nCO_2H$, $-(CH_2)_nCONH_2$, $-(CH)_n$ phenyl, $-(CH)_n$ phenylalcohol, $-(CH_2)_n$ indole, $-(CH_2)_n$ imidazole, $-(CH_2)_xOH$, $-(CH_2)_xSH$, $-(CH)_xS$ C_1 - C_6 alkyl, $-(CH_2)_xNHC(NH)NH_2$ or $-(CH_2)_xNH$ in which x is 1 to 5 and n is 0 to 5; or **R3** and **R4** are together taken with C_1 - C_4 to form a ring; **R5** represents NR_6R_7 , $-OR_8$ or $-OCH(R_8)OCOR_9$ in which R_6 and R_7 are each independently hydrogen, C_1 - C_6 alkyl or phenyl substituted with C_1 - C_4 alkoxy, nitro or halogen and R_8 and R_9 are each independently C_1 - C_8 alkyl, C_3 - C_{11} cycloalkyl substituted with halogen, nitro, hydroxy, C_1 - C_4 alkyl or C_1 - C_4 alkoxy or phenyl substituted with halogen, nitro, hydroxy, C_1 - C_4 alkyl or C_1 - C_4 alkoxy; and **W** represents oxygen or sulfur, which comprises reacting a compound of the general formula II:



Formula II

in which **R1**, **R2** and **W** represent the same as defined above, with a compound of the general formula III:



Formula III

10 in which R3, R4 and R5 represent the same as defined above, to produce the said compound of the general formula I.

6. A pharmaceutical composition useful in the inhibition of phosphodiesterase IV or tumor necrosis factor comprising pharmaceutically effective amount of the compound according to claim 1 and a pharmaceutically acceptable carrier.

15

20

25

30

INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR 97/00156

A. CLASSIFICATION OF SUBJECT MATTER

IPC⁶: C 07 C 235/52,237/22; C 07 D 207/16; C 07 C 323/02,231/02; A 61 K 31/22, 31/165
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC⁶: C 07 C 253/52,237/22,323/59,231/02; C 07 D 207/16; A 61 K 31/22,31/165

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Chemical Abstracts, Vol.105, No.11, 15 September 1986 (Columbus, Ohio, USA), page 88, column 1, abstract No.91509y, MAKOVEC, F. et al.: "New glutamic and aspartic derivatives with potent CCK-antagonistic activity", & Eur. J. Med. Chem.--Chim. Ther. 1986, 21(1), 9-20 (Eng). Chemical Abstracts, Eleventh Collective Index, Vol.96-105, FORMULAS, page 14736F, C20H3ON2O6, Pentanoic acid, 4-[(3,4-dimethoxybenzoyl)amino]-5-(dipropylamino)-5-oxo (=).	1
X	Chemical Abstracts, Vol.114, No.11, 18 March 1991 (Columbus, Ohio, USA), page 14, column 2, abstract No.94611x, COATS, E.A. et al.: "Correlation analysis and molecular modeling of cholecystokinin inhibitors", & Quant. Struct.-Act. Relat. 1990, 9(2), 94-101 (Eng). Chemical Abstracts, Twelfth Collective Index, Vol.106-115, FORMULAS, page 20045F, C20H3ON2O6, Pentanoic acid, 4-[(3,4-dimethoxybenzoyl)amino]-5-(dipropylamino)-5-oxo-(S).	1,3

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

23 January 1998 (23.01.98)

Date of mailing of the international search report

05 February 1998 (05.02.98)

Name and mailing address of the ISA/AT
 AUSTRIAN PATENT OFFICE
 Kohlmarkt 8-10
 A-1014 Vienna
 Facsimile No. 1/53424/535

Authorized officer

Körber

Telephone No. 1/53424/457

INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR 97/00156

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Chemical Abstracts, Vol.123, No.17, 23 October 1995 (Columbus, Ohio, USA), page 853, column 2, abstract No.313505g, PACE, R.D. et al.: "Allylboration of α -Amino Ketones", & J. Org. Chem. 1995, 60(15), 4838-44 (Eng). Chemical Abstracts, Vol.123, Formula Index, page 3109F, C19H21N05, L-Phenylalanine, N-(3,4-dimethoxybenzoyl)-methyl ester.	1,3
X	Chemical Abstracts, Vol.123, No.17, 23 October 1995 (Columbus, Ohio, USA), page 1190, column 1, abstract No.228176f, OKURA, A. et al.: "Preparation of heterocyclic compounds as tyrosine kinase inhibitors", & WO 95/06 032 (BANYU PHARMACEUTICAL CO., LTD.). Chemical Abstracts, Vol.123, Formula Index, page 1943F, C13H17N05, Glycine, N-(3,4-dimethoxybenzoyl)-ethyl ester.	1
X	Chemical Abstracts, Vol.99, No.23, 05 December 1983 (Columbus, Ohio, USA), page 738, column 2, abstract No.194796b, KNEFELI, F. et al.: "Electron-impact induced loss of C-5 and C-8 substituents in 1,2,3,4-tetrahydroisoquinolines. I. Synthesis of 4-acetylpyrrolo 1,2-b isoquinoline", & Arch. Pharm. (Weinheim, Ger.) 1983, 316(9), 773-81 (Ger). Chemical Abstracts, Eleventh Collective Index, Volumes 96-105, Formulas, page 10025F, C15H19N05, L-Prolin, 1-(3,4-dimethoxybenzoyl)methyl ester.	1,3
X	US 2 956 081 A (G.W. KUSSEROW) 11 October 1960 (11.10.60), column 1, line 55 - column 2, line 4; example 6.	1,5
X	EP 0 388 967 A1 (WAKAMOTO PHARMACEUTICAL CO.)	1
Y	26 September 1990 (26.09.90), page 14, line 44.	5
Y	WO 97/05 105 A1 (PFIZER INC.) 13 February 1997	5
A	(13.02.97), page 1, lines 8-36; page 7, lines 14-24.	1-3,6
A	WO 93/15 044 A1 (SMITH-KLINE BEECHAM CORPORATION) 05 August 1993 (05.08.93), pages 1,2.	6

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.

PCT/KR 97/00156

Im Recherchenbericht angeführtes Patentdokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication
US A 2956081		keine - none - rien	
EP A1 388967	26-09-90	DE C0 69020394 DE T2 69020394 EP B1 388967 HU A0 901814 HU A2 57744 HU A3 9500234 HU B3 211520 US A 5055481 US E 35281 JP A2 4270271	03-08-95 29-02-96 28-06-95 28-07-90 30-12-91 28-09-95 28-11-95 08-10-91 18-06-96 25-09-92
WD 9705105		keine - none - rien	
WD A1 9315044	05-08-93	AU A1 35886/93	01-09-93